

Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at http://about.jstor.org/participate-jstor/individuals/early-journal-content.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

"I gave my son directions to show you two other coins of the same set, which you possibly will also reckon Etruscan.

"I am, dear Sir William,
"Your faithful Servant,
"THOMAS L. COOKE."

The following notice was communicated by Sir William Rowan Hamilton, of a Paper "on the Application of Quaternions to the Determination of the Distance of any recently discovered Comet or Planet from the Earth."

This celebrated problem is treated in this paper by means of the formulæ which were communicated to the Academy by the author, in July, 1845. The chief step consists in a very easy deduction, from those formulæ, of the equation:

$$\frac{a}{c}\left(\frac{M}{a^3} - \frac{M}{b^3}\right) = \frac{S \cdot \gamma \gamma' \gamma''}{S \cdot \gamma \gamma' a};$$

where c is the sought distance of the comet (or planet) from the earth; M is the mass of the sun, and a and b are the distances of earth and comet from that body; a is the heliocentric vector-unit of the earth, and γ is the geocentric vector-unit of the comet; while γ' , γ'' are the first and second differential coefficients of γ , taken with respect to the time, and determined, along with γ itself, from three successive observations: and S is the characteristic of the operation of taking the scalar part of a quaternion. The second member of the equation admits of being geometrically interpreted as a ratio of two pyramids, and can in various ways be transformed by the rules of the calculus of quaternions.

Mr. Donovan exhibited a table gas lamp of his invention, which generates its own gas, and made the following statement relative to it: